Structure of the Earth: Mantle and Core

THORNE LAY

*Department of Geological Sciences
University of Michigan
Ann Arbor, MI 48109*

Abstract. The deep interior structure of the earth has been extensively analyzed using a wide variety of seismic phases and techniques during the last four years. Most studies have emphasized quantitative three-dimensional mapping of the lateral velocity heterogeneity of the mantle and core. These aspherical velocity variations are believed to be direct manifestations of thermal and compositional heterogeneity associated with convective processes. A first generation of global models for the lateral velocity variations in the deep earth has been produced, providing tantalizing images of large scale structures suggestive of a non-steady state thermal convection system.

The upper 400 km of the mantle has the strongest lateral velocity variations, of up to $\pm 10\%$ for shear velocity. Surface-wave analyses that do not require *a priori* regionalizations have demonstrated that there is a strong association between surface tectonic provinces and the uppermost mantle velocity variations. Thus, the thermal and convective state of the upper 200 km of the mantle can be reliably interpreted in the context of plate tectonics. The upper mantle models support the contention that continents have deep roots, with differences in velocity structure from oceanic and active tectonic regions extending as deep as 400 km. Very long-wavelength lateral velocity variations of a few percent have been detected in the transition zone at depths from 400 to 670 km, as well as throughout the lower mantle. These deep-seated variations have little correspondence to surface tectonics, and efforts to interpret their nature are just beginning. The lowermost 200 km of the mantle (D" region) has lateral velocity fluctuations comparable to those in the upper mantle, and evidence has been presented for the presence of a sizable velocity discontinuity at the top of the D" layer. A combined thermal and compositional boundary layer, roughly mirroring the lithosphere, is a likely explanation for this anomalous zone. The core-mantle boundary appears to have significant (10 km) long-wavelength topography, presumably sustained by dynamic stresses from deep mantle convection. The inner core may have strong lateral heterogeneity or axially symmetric anisotropy, suggesting a complex thermal and compositional state.

Significant progress has been made in characterizing the frequency dependence of anelastic attenuation in the mantle in the short-period body-wave band. Models for teleseismic P-wave attenuation operators have converged, with t^* values of 0.7-1.0 s appropriate at 1 Hz, and t^* values of 0.4-0.6 s appropriate at 4 Hz. Regional variations of attenuation are slowly being mapped out as well.

MANTLE STRUCTURE

An unprecedented increase in our knowledge of mantle structure has been achieved in the last four years. A broad spectrum

of seismological techniques, including body-waveform modeling, global travel-time inversion, surface-wave dispersion and waveform inversion, and normal-mode splitting analysis, have been brought to bear upon the problem of mapping the velocity deviations from radially symmetric earth structure in the deep interior. While this lateral velocity heterogeneity is relatively small, typically being only a few percent variation about the average earth model, its significance is profound. This is because the velocity variations are signatures of thermal, compositional, and crystal orientation differences, all of which are associated with mantle dynamics. Accurate three-dimensional imaging of the heterogeneity thereby provides the most direct means for determining the configuration of mantle convection systems. The seismological models also serve as critical constraints for geodynamic and geochemical modeling of the earth, and are resulting in increasing interdisciplinary interaction. Perhaps the most important inference to be drawn from the first generation of global models is that the rich complexity of the interior indicates a non-steady system, for which simplistic notions of steady-state flow may be inapplicable.

Surface-Wave Tomography

The accumulation of a substantial data base of long-period digital recordings from Global Digital Seismic Network (GDSN) and International Deployment of Accelerometers (IDA) instruments has enabled, for the first time, systematic analyses of surface-wave dispersion on a global scale. Additional large data bases of body-wave travel-times collected by the ISC or from special readings of WWSSN recordings have enabled corresponding global and continental scale body-wave investigations. The process of inverting these large sets of path anomalies for a heterogeneous model has been called seismic tomography [Anderson and Dziewonski, 1984; Dziewonski and Anderson, 1984]. Several different procedures have been used to analyze the surface-wave observations. One procedure involves the standard analysis of isolated fundamental-mode wavetrains to extract phase and group velocities, which are subsequently inverted for upper mantle structure. Another procedure involves direct waveform modeling of the entire seismic trace, including overtones and amplitudes. The latter procedure directly results in a velocity structure, but must be performed iteratively, and has so far been implemented only with asymptotic approximations to the normal-mode theory of Woodhouse and Girnius [1982] that make it almost equivalent to the step by step dispersion analysis.

Nakanishi and Anderson [1982, 1983, 1984a,b] adopted the approach of inverting the dispersion measurements for spherical harmonic expansions of Love- and Rayleigh-wave phase and group velocity up to degree and order 6. By measuring both odd and even orbit arrivals at single stations, the odd harmonics of the expansion can be determined, albeit with greater uncertainty than for the even harmonics, which are constrained by more accurate great-circle measurements. The spherical harmonic

Copyright 1987 by the American Geophysical Union.

Paper number 6R0768.
8755-1209/87/006R-0768\$15.00

expansions do not require an *a priori* tectonic regionalization. Nataf et al. [1984, 1986] inverted the dispersion measurements for transversely isotropic models that explicitly include shear-wave and compressional-wave anisotropy. Other inversions of the dispersion data were performed by Tanimoto [1985, 1986b] and Tanimoto and Anderson [1984, 1985]. The analyses by Tanimoto utilized Backus-Gilbert inverse theory to appraise the issue of error and resolution of the phase velocity models and associated shear velocity inversions. The error analysis illustrated the significance of the antipodal peak in the resolution kernel, which, in combination with statistical errors, seriously limits the resolvable anomalies. The resolution kernel for the data sets of 400 (Love) and 600 (Rayleigh) phase velocity measurements used in these studies is localized within an area of 2000 km radius. Limited azimuthal path coverage of most areas of the earth prevents inversions for global azimuthal anisotropy at present [Tanimoto and Anderson, 1985], and differences in the depth resolution of the Love (SH) and Rayleigh (P, SV) modes leaves open the question of how well transverse isotropy below 200 km can be resolved with the current data [Tanimoto, 1986b].

Woodhouse and Dziewonski [1984] and Woodhouse [1984] used a much larger data set of waveforms from 53 earthquakes and 870 paths in a complete waveform inversion procedure. The inversion resulted in a global shear-wave velocity model with a spherical harmonic expansion up to degree and order 8, and a cubic polynomial description with depth for the upper 670 km of the mantle. Their analysis was restricted to signals with periods longer than about 135 s, close to the 150 s cutoff of the dispersion analyses.

The very long-wavelength features in the models obtained from the dispersion and waveform procedures are remarkably similar, suggesting that first-order characteristics of the actual mantle structure have been resolved. All of the inversions have demonstrated that the strongest heterogeneities are in the upper 250 km of the mantle, and these have a strong correlation with surface tectonic features. Ocean ridges and west Pacific subduction zones are underlain by slow velocities, while all major shields have fast velocity roots, some of which extend to at least 350 km depth. The South Atlantic has relatively high velocities at shallow depths, and low velocities under the Red Sea and Gulf of California persist to 400 km depth. Below 250 km many portions of ridges are underlain by high velocity regions, notably the South East Indian Ridge. Many of these features would not have been anticipated in *a priori* regionalized models. The anisotropic inversions of Nataf et al. [1986] suggest a belt of SV>SH anisotropic velocities at depths of 200 to 400 km following the circum-Pacific ridge and subduction systems, suggesting vertical flow; however, similar features were not resolved by Tanimoto [1986b]. The procedure used by Woodhouse and Dziewonski [1984] appears to resolve mantle structure at somewhat greater depths due to the inclusion of overtone data, and their model shows a strong degree-2 component in the transition zone (400-670 km) with broad regions of high velocity under South America and the South Atlantic and under the western Pacific, and broad regions of low velocities under the central and eastern Pacific and under the Middle East. The velocity anomalies in these models are on the order of $\pm 8\%$ at 50 km depth, $\pm 2.5\%$ at 450 km, and $\pm 1\%$ at 650 km.

The global models indicate, but have limited resolution of fine features such as the evolution of velocity structure of oceanic lithosphere with age. Greater resolution of oceanic structure,

particularly for the Pacific Ocean, has been obtained in the regionalized dispersion studies of Anderson and Regan [1983], Regan and Anderson [1984], Chao et al. [1983], Dziewonski and Steim [1982], Knopoff [1982], Nishimura and Forsyth [1985, 1986], and Rosa and Aki [1986]. Detailed continental dispersion analysis of the crust and upper mantle under Eurasia was performed by Feng and Teng [1983]. Improving the spatial resolution of both the global and smaller scale inversions requires the use of shorter period data, but the great-circle path assumption employed in the dispersion and waveform modeling procedures has dubious validity for periods less than 150 s. Several studies have addressed the great-circle assumption using surface-wave raytracing [Lay and Kanamori, 1985; Schwartz and Lay, 1985; Tajima and Garmany, 1986; Wong and Woodhouse, 1983], and by appraising the effects of heterogeneity on moment tensor inversions [Nakanishi and Kanamori, 1982; Tanimoto and Kanamori, 1986]. The recognition that strong amplitude and phase effects due to departure from great-circle paths can occur for periods as long as 300 s has led to efforts to directly utilize the focussing and defocussing information in inversions for the heterogeneous models [Wong and Woodhouse, 1986; Yomogida, 1985; Yomogida and Aki, 1985, 1986]. In a parallel development, Tanimoto [1984] obtained degree-2 Love-wave models using a Born approximation in a waveform inversion procedure that explicitly utilized long-period amplitude information.

Other surface-wave procedures for improving the resolution of the upper mantle models have involved detailed analysis of overtones, which sample the transition zone more completely than the fundamental-modes. Lerner-Lam and Jordan [1983] extracted the higher-mode information using branch cross-correlation functions between single-mode branch synthetics and the observed seismograms. This procedure was used to develop upper mantle shear velocity structures across Eurasia and the eastern Pacific. The resolution capability of this procedure was analyzed by Gee and Jordan [1986], and an extension to anisotropic inversion was considered by Lerner-Lam [1986]. Okal and Jo [1983, 1985] have analyzed dispersion characteristics of longer period spheroidal overtones. One of the most promising procedures appears to be inclusion of body-wave and early overtone signals in the global surface-wave inversions, as shown by Woodhouse and Dziewonski [1986]. Further progress will be possible as instruments which remain linear for the first surface-wave arrivals from large earthquakes are deployed.

Free-Oscillations

The surface-wave investigations described above have proven most fruitful in leading to models of upper mantle heterogeneity because of asymptotic approximations for the traveling waves which lead to straightforward data analysis and inversion. Analysis of the splitting of normal-mode spectra due to ellipticity, rotation and lateral heterogeneity is also beginning to yield aspherical earth structure models, and formalisms are being developed that should eventually lead to non-asymptotic inversions.

Masters et al. [1982] inverted multiplet locations for the degree-2 spherical harmonic component of the earth's heterogeneity. Their preferred model localized the heterogeneity in the transition zone, with shear velocity variations on the order of 1.7%. This interpretation was questioned by Kawakatsu [1983], who showed that some regionalized models of shallow hetero-

geneity have a correspondingly strong degree-2 component, but the transition zone feature was also obtained in the surface-wave inversion by Woodhouse and Dziewonski [1984]. Giardini et al. [1986] have recently inverted split-mode data for mantle heterogeneity models, finding that small perturbations about the surface-wave derived models can explain the mode data that are sensitive to the upper mantle.

Other free-oscillation research has concentrated on mode-coupling considerations, which have proved more significant than was previously recognized, and on procedures for calculating the normal-modes for an aspherical earth model. Tanimoto and Bolt [1983] analyzed coupling of toroidal-modes, and Masters et al. [1983] presented observations of coupling between spheroidal- and toroidal-modes at very low frequencies. Theoretical treatments of the aspherical earth calculations have been presented by Dahlen and Henson [1985], Morris and Geller [1982], Park [1986], Park and Gilbert [1986], Tanimoto [1984], and Tsuboi et al. [1985]. Davis [1985] has considered variations in apparent attenuation resulting from lateral heterogeneity, and Davis and Henson [1986] addressed the validity of the great-circle assumption for normal-mode measurements.

Efforts to refine radially symmetric mantle models and mode measurement procedures have been presented by Davis [1986], Henson [1982] and Masters and Gilbert [1985]. The effects of anisotropy on the normal-modes has been considered by Anderson and Dziewonski [1982] and Tanimoto [1986a]. The development of normal-mode partial derivatives for general anisotropy will be important for the next generation of upper mantle surface-wave and normal-mode inversions for anisotropic velocity structure.

Body-Waves

A variety of body-wave phases have been analyzed to determine both radial and lateral velocity structure in the upper mantle. WWSSN recordings of long-period transverse component S and SS phases have been particularly fruitful. A series of waveform modeling studies of these phases by Given [1983], Grand and Helmberger [1984a,b, 1985], Graves et al. [1985], Helmberger et al. [1985a,b], and Rial et al. [1984] has led to detailed shear velocity models for tectonic regions, shield regions, young and old oceanic provinces, and the transitional zones connecting tectonic provinces. These studies have not only refined our knowledge of the size of upper mantle discontinuities and of the velocity gradients between them but have also established that continental roots persist to 400 km depth and that 10% lateral variations in shear velocity exist near 200 km depth. Travel-time anomalies for the SS reverberations have been used to develop continental scale images of the upper mantle heterogeneity beneath North America [Grand, 1986a,b] the Atlantic [Kuo et al., 1986], and the Indian Ocean [Stark and Forsyth, 1983], with a resolution that is not viable with the global surface-wave inversions. Corresponding PP-wave analysis has just begun [Lefevre and Helmberger, 1984].

Analysis of short-period P-waves recorded by the large Southern California array has resulted in models of the lateral heterogeneity under California, as well as radial P-wave models for the upper mantle structure of a tectonically active region. Humphreys et al. [1984] used tomographic inversion to refine the image of a high velocity anomaly in the upper 300 km beneath the Transverse Ranges. Walck [1984, 1985] and Walck and Clayton [1984] analyzed a dense array of data from the upper mantle triplication range to develop radial P-wave

velocity models, placing tight constraints on the upper mantle velocity gradients and size of the discontinuities at 400 and 650 km depth. These models, together with the S-wave structures from the SS analyses have been used to bound petrological models for the upper mantle by Anderson and Bass [1984], Bass and Anderson [1984], and Jeanloz and Thompson [1983]. Underside reflections of P'dP' have also been analyzed to appraise the sharpness of the discontinuities [Lees et al. 1983; Murtha and Tanimoto, 1982], as have long period ScS-type reverberations [Revenaugh and Jordan, 1985].

An increasing number of studies are attempting to extract upper mantle structure from SV component recordings, which have always been more difficult to interpret than the SH signals. Baag and Langston [1985a,b, 1986], Baumgardt and Alexander [1984], Langston and Baag [1985], and Zandt and Randall [1985] have performed modeling calculations for Sp and SPL phases that are particularly sensitive to uppermost mantle structure. Propagation of S-waves in heterogeneous media has been considered by Cormier [1984, 1986], who appraised the effects on the S-wave polarization. Observations of S-wave splitting attributed to anisotropy were presented by Ando et al. [1984].

Body-wave travel-time and amplitude station anomalies have been used to investigate upper mantle heterogeneity on a variety of scales. Taylor [1983], Lynnes and Lay [1984], and Priestley and Chavez [1985] have considered the travel-time and focusing effects of upper mantle heterogeneity beneath the Nevada Test Site, which appears to be related to deep roots of the caldera system in the region. Michaelson and Weaver [1986] analyzed upper mantle structure under the Pacific Northwest using travel-time delays. Variations of P-wave amplitudes and travel-times across the North American continent have been considered by Butler [1983, 1984a,b, 1985] and Lay and Helmberger [1983a], while global P-wave [Dziewonski and Anderson, 1983] and S-wave [Souriau and Woodhouse, 1985] travel-time anomalies have been presented and related to temperature and compositional variations.

Slab Structure

Subducted oceanic lithosphere constitutes the strongest localized velocity heterogeneity in the upper mantle. Numerous seismicity and body-wave travel-time studies have addressed subducting slab geometry and velocity structure. Complexity of the stress state in subducted slabs has been revealed by studies of the multiple Benioff zones found in some slabs [Kawakatsu, 1985, 1986a,b; Kawakatsu and Seno, 1983; Cahill and Isacks, 1986; Taber and Hudnut, 1984], as well as by studies of focal mechanisms and seismicity levels [Giardini and Woodhouse, 1984; Vassiliou, 1984; Vassiliou et al., 1984]. The configuration of various subducted slabs, as indicated by seismicity and velocity heterogeneity, has been analyzed by Boyd et al. [1984], Chiu et al. [1985], Cockerham [1984], Grange et al. [1984], Hauksson [1985], Roecker [1985], and Rohay [1982]. Several studies have addressed the effects of slab heterogeneities on propagating waves as well [Bolt and Drake, 1986; Langston and Arnold, 1982].

The most significant advances in our knowledge of slab structure concern the depth of penetration and sharpness of the velocity gradients of the slab. Creager [1984] and Creager and Jordan [1984, 1986a] analyzed travel-time residual patterns from intermediate and deep focus earthquakes in western Pacific subduction zones, finding systematic patterns that can

be well explained by thermal models of subducting slabs that penetrate into the lower mantle to depths of at least 1000 km, well below the depth of the deepest earthquakes [Stark and Frohlich, 1985]. Similar analyses of travel-time residual patterns for events in Tonga [Fischer et al., 1986] and the Aleutians [Spencer and Engdahl, 1983; Boyd and Creager, 1986] also indicate aseismic extensions of the slab velocity anomaly below the maximum depth of seismicity. Other investigations of the sharpness of the velocity gradients in the slab [Stefani et al., 1982]; the velocity structure in the mantle surrounding the slab [Suyehiro and Sacks, 1985]; and anisotropy of the slab itself [Anderson, 1986], have not refuted the lower mantle slab penetration hypothesis. Silver and Chan [1986] and Beck and Lay [1986] have analyzed waveform complexity in broadband S waves that appears to arise from interaction of the waves with strong velocity heterogeneity in the lower mantle. Azimuthal patterns and travel-time correlations in these data are consistent with multipathing in a lower mantle slab extension. Although further work on the slab penetration hypothesis is clearly needed, it presently appears that a strong case has been made for mass flux across the 650 km discontinuity, which has profound consequences on the issue of whole mantle versus layered convection.

LOWER MANTLE STRUCTURE

In addition to the slab structures that appear to extend into the lower mantle, velocity heterogeneity at a variety of scales has been detected below the 670-km discontinuity. Surface-wave analysis cannot be used to determine the structure in the lower mantle, but body-wave travel-times reveal both small and large scale variations. Localized velocity heterogeneities with scale lengths of about 1000 km and 2% velocity heterogeneity were detected beneath the Caribbean and South America by Lay [1983], using S and ScS travel times from WWSSN recordings. A high velocity anomaly beneath the Caribbean at depths of 800 to 1900 km was found to extend northward, with a systematic offset to the east with increasing depth, in the tomographic inversion by Grand [1986a]. A similar, elongate high velocity ridge beneath Alaska is apparent in the recent inversions of Woodhouse and Dziewonski [1986], which incorporated body-wave data in the global inversion. The geodynamic significance of these structures is not yet established.

Several studies have used the vast travel-time data base accumulated by the ISC to investigate lower mantle structure. Lee and Johnson [1984a,b] established extremal bounds on radial velocity models for the lower mantle using Tau-p measurements from the ISC travel times. Dziewonski [1984] used a time-term approach to analyze about 500,000 travel-time residuals from 5000 earthquakes in deriving an aspherical P velocity model for the lower mantle. The model was represented by spherical harmonics up to degree 6 and radial order number 4. The earthquakes were relocated in each iteration for the structure, and weighting schemes were used to suppress the aliasing due to non-uniform coverage of the interior. The model has strong velocity perturbations of 1-1.5% just below the 670-km discontinuity and just above the core-mantle boundary. At a depth of 2000 km the perturbations in velocity are about 3 to 4 times smaller than at the two boundaries. Qualitatively similar results were obtained in the ISC travel-time inversions of Clayton and Comer [1983] and Comer and Clayton [1986], who used a tomographic procedure with blocks rather than an analytic representation. This procedure provides higher spatial

resolution, and can easily be extended to include travel-times from phases other than P [Zhou and Clayton, 1985]. The low order variations of the two approaches are quite similar, and have been shown to successfully predict long-wavelength components of the geoid under the assumption of a dynamic, viscous earth by Hager et al. [1985], who applied the theory of Hager [1984] and Richards and Hager [1984]. The large-scale velocity variations apparent in these models do not have any simple relation to surface tectonics, and appear to have much greater complexity than anticipated for a simple, steady-state convection system. Two possible interpretations are that there are significant compositional heterogeneities entrained in the flow [Davies, 1984], or alternatively, the flow is not steady-state at all.

Structure near the Core-Mantle Boundary

While the global tomographic inversions can resolve the presence of large-scale heterogeneity near the base of the mantle, other procedures are required to model the detailed structure. Accurate velocity models are needed in order to assess recent thermal calculations that predict a major, unstable thermal boundary layer at the base of the mantle that serves as a source of thermal plumes [Boss and Sacks, 1985; Loper, 1984; Loper and Stacey, 1983; Stacey and Loper, 1983]. Short-period P-waves near the onset of the core shadow zone indicate substantial lateral variation in average radial structure of the lowermost 200 km of the mantle (D'' region) [Ruff and Lettvin, 1986]. Model experiments have been conducted by Menke [1986a,b] to assess the behavior of short period PcP phases. PKP phases have also been analyzed, and appear to require short scale length heterogeneities in D'' [Snoke and Sacks, 1986]. Strong heterogeneity in the S-wave velocities of D'' are indicated by the diffracted S-wave study of Bolt and Niazi [1984] and the ScS travel time study of Lavelle et al. [1986].

A series of modeling studies of long period SH and SV signals traversing D'' by Lay [1985, 1986a,b], Lay and Helmberger [1983b,c,d], Lay and Young [1986], Young and Lay [1986a], and Zhang and Lay [1984] indicate the presence of waveform complexities that can be well-modeled by shear velocity models with a 2.75% shear discontinuity about 280 km above the core-mantle boundary. A reflection from this velocity increase is apparent in transverse component seismogram profiles at distances greater than 70°, with data sampling four different regions of D'' having consistent waveform characteristics. Waveform modeling of long-period ScSH and ScSV indicates the presence of either fine structure just above the core or possibly anisotropy in D'' [Lay and Helmberger, 1983d; Cormier, 1985]. At this time the seismic models for D'' are not consistent enough to resolve the nature of the thermal boundary layer, as is discussed in the review by Young and Lay [1986b], and it appears that the presence of both a thermal and a chemical boundary layer at the base of the mantle is the best way to explain the radial and lateral structure of the region [Jordan and Creager, 1986].

CORE STRUCTURE

The radial and lateral velocity structure of the core has received increasing attention, particularly with the availability of models of the mantle heterogeneity that allow the shallow travel-time effects to be removed from data that sample the core. Large ISC travel-time data sets have been used to determine extremal bounds on the radial velocity models for the core

[Johnson and Lee, 1985; Stark et al., 1986]. Detailed analysis of SKS travel-times at distances less than 95° provide control on the velocity gradient in the outermost core [Murtha, 1984; Lay and Helmberger, 1983d], which appears not to have an anomalously steep value. Waveform modeling studies of broadband PKIKP phases indicate a simple inner core-outer core discontinuity [Choy and Cormier, 1983; Cormier and Choy, 1986], consistent with the radial travel time inversions.

However, the first generation of three-dimensional core models indicate greater complexity of core structure. Jordan and Creager [1986] and Morelli et al. [1986] have used large data sets of PKP and PKIKP travel times from the ISC bulletins to image the low-order heterogeneity of the core. Both studies indicate long-wavelength heterogeneity that appears to reside in both the inner core and near the core-mantle boundary. Topography on the core-mantle boundary of ± 8 km or so with very long wavelengths can account for much of the anomaly in PKP times. Some of the heterogeneity may actually be in the outer core, but dynamic considerations prevent any density heterogeneity from persisting in this region. Anomalously large splitting of normal-modes sensitive to the core structure also indicates very long-wavelength aspherical structure. Ritzwoller et al. [1986] concluded that an axisymmetric structural anomaly in the outer core is required to explain these observations. Giardini et al. [1986] developed an inversion procedure utilizing the anomalously split modes and derived core structures with 8 km undulations of the core-mantle boundary, 25 km undulations of the inner core-outer core boundary, and several percent velocity variations within the inner core, dominated by zonal harmonics. Morelli et al. [1986] and Woodhouse et al. [1986] have proposed an alternate model in which the inner core has a strong axially symmetric anisotropy, which produces systematic PKIKP anomalies and anomalous mode splitting. The complexity of the inner core appears compatible with the models for this boundary proposed by Loper and Fearn [1983] based on geodynamic and geochemical arguments.

ANELASTIC PROPERTIES OF THE DEEP EARTH

With the exceptions of the normal-mode attenuation study by Masters and Gilbert [1983], the overtone attenuation study of Okal [1986] and the surface-wave attenuation study of Patton and Taylor [1984], most seismological research on attenuation in the last four years has involved the body-wave frequency band. Progress in this area has been substantial, largely due to the absorption band model concepts that were developed several years ago [see review by Cormier, 1982]. The frequency dependence of short-period P-wave attenuation and its regional variations have been studied using spectral shape measurements by Bache [1985], Bache et al. [1985, 1986], Der and Lees [1985], Der et al. [1982a,b, 1985, 1986] and Shore [1984]. Broadband P and S data have been analyzed to infer regional variations across North America by Taylor et al. [1986]. These studies have clearly demonstrated the decrease in attenuation for frequencies greater than 1 Hz. Additional studies have reliably established the absolute attenuation levels by comparing near-field source models with teleseismic observations [Burdick et al., 1984; Burger et al., 1986], or by source cancellation experiments using phases such as ScS-ScP [Burdick, 1985], sS-sP [Burdick and Grand, 1984] and multiple ScS phases [Lay and Wallace, 1983]. Consistent frequency dependent models for short-period P-waves have resulted from the different studies, with t^* values of 0.7 to 1.0 s for 1 Hz signals and 0.4 to 0.6 s for 4 Hz signals.

Deep mantle attenuation studies using PcP phases [Bolt and Canas, 1985] and ScS phases [Choy and Cormier, 1986] do not appear to support early models with a very low Q zone at the base of the mantle, although the recent model of Shore [1984] has such a feature in it. The effects of diffraction and the increased velocity heterogeneity in D'' need to be accounted for when producing Q models for the lowermost mantle. The apparent attenuation effects of scattering have received some attention [Menke, 1983; Richards and Menke, 1983], though it has proven difficult to separate intrinsic attenuation from scattering losses in actual data.

CONCLUSIONS

The large number of deep earth structure studies reviewed above reflects a recent emphasis on quantitative mapping of the three-dimensional velocity structure of the earth's interior. This has long been a principal goal of seismology, but it is only recently that high quality seismic data bases have been available for global analysis. The excitement generated by the first three-dimensional models of the aspherical heterogeneity has breached disciplinary boundaries and has given strong impetus to the recent efforts to upgrade the global network of seismometers [IRIS, 1984]. The rapid progress in imaging the interior using both digital and WWSSN data is a testimonial to the benefits to be reaped from high quality global network data.

It appears that the earth is heterogeneous at all depths and at all scales; thus it will be a difficult task to accurately determine the detailed structure of the interior. Error and resolution analysis will be increasingly important for appraising the results of large inversion procedures, which are often subject to subtle biases and instabilities. An open mind with regard to the nature of the earth's structure, particularly regarding general anisotropy, boundary layer structure, and whole mantle convection is important in these modeling efforts. The development of aspherical earth models has been paralleled by development of techniques for modeling wave propagation in heterogeneous media, and inverse procedures will have to be developed to exploit these.

Studies of the core-mantle boundary and core structure have begun to reveal unexpected complexity that has important geodynamic and geochemical consequences. It will be particularly important to integrate body-wave and free-oscillation data bases to better constrain the heterogeneity of this region. Establishing the role of the core-mantle boundary as either a thermal or a chemical boundary layer is particularly important for the geothermal models of the interior.

The question of lower mantle slab penetration has not been completely resolved, and quantitative three-dimensional wave-field modeling is needed to further test the travel-time models. Additional studies of the 670-km discontinuity, especially in the vicinity of subducted slabs are needed to establish the nature of this major boundary. Similarly, additional studies of the nature of upper mantle transitions between shield and tectonic provinces are needed to better resolve whether thermal, chemical, or anisotropic variations are responsible for the strong contrasts.

Acknowledgements. This research was partially supported by NSF grant EAR-8451715, a grant from the Alfred P. Sloan Foundation, and a Shell Faculty Career Initiation Grant. Acknowledgment is made to the Donors of the Petroleum Research Fund administered by the American Chemical Society for the partial support of this research.

REFERENCES

- Anderson, D. L., A new look at the inner core of the earth, *Nature*, 302, 660, 1983.
- Anderson, D. L., Surface wave tomography, *Eos Trans. AGU*, 65, 147-148, 1984.
- Anderson, D. L., Evolution of earth structure and future directions of 3D modeling, in *The VELA Program*, ed Ann Kerr, 399-418, 1985.
- Anderson, D. L., The mineralogy of deep slabs, *Eos Trans. AGU*, 67, 379, 1986.
- Anderson, D. L., and J. D. Bass, Mineralogy and composition of the upper mantle, *Geophys. Res. Lett.*, 11, 637-640, 1984.
- Anderson, D. L., and A. M. Dziewonski, Upper mantle anisotropy evidence from free oscillations, *Geophys. J. R. Astron. Soc.*, 69, 383-404, 1982.
- Anderson, D. L., and A. M. Dziewonski, Seismic tomography, *Scientific American*, 251, 60-68, 1984.
- Anderson, D. L., and J. Regan, Upper mantle anisotropy and the oceanic lithosphere, *Geophys. Res. Lett.*, 10, 841-844, 1983.
- Ando, M., Y. Ishikawa, and F. Yamazaki, Shear wave polarization anisotropy in the upper mantle beneath Honshu, Japan, *J. Geophys. Res.*, 88, 5850-5864, 1984.
- Baag, C.-E., and C. A. Langston, Shear-coupled PL, *Geophys. J. R. Astron. Soc.*, 80, 363-385, 1985a.
- Baag, C.-E., and C. A. Langston, A WKBJ spectral method for computation of SV synthetic seismograms in a cylindrically symmetric medium, *Geophys. J. R. Astron. Soc.*, 80, 387-417, 1985b.
- Baag, C.-E., and C. A. Langston, Diffracted Sp generated under the Australian Shield, *J. Geophys. Res.*, 91, 9507-9516, 1986.
- Bache, T. C., The effect of Q on teleseismic P waves, in *The VELA Program*, ed Ann Kerr, 252-272, 1985.
- Bache, T. C., S. R. Bratt, and H. Bungum, High frequency P wave attenuation along five teleseismic paths from central Asia, *Geophys. J. R. Astron. Soc.*, 85, 505-522, 1986.
- Bache, T. C., P. D. Marshall, and L. B. Bache, Q for teleseismic P waves from central Asia, *J. Geophys. Res.*, 90, 3575-3587, 1985.
- Bass, J. D., and D. L. Anderson, Composition of the upper mantle: geophysical tests of two petrological models, *Geophys. Res. Lett.*, 11, 237-240, 1984.
- Baumgardner, D. R., and S. S. Alexander, Structure of the mantle beneath Montana: LASS analysis of long-period, mode converted phases, *Bull. Seismol. Soc. Am.*, 74, 1683-1702, 1984.
- Beck, S. L., and T. Lay, A test of the lower mantle slab penetration hypothesis using broadband S waves, *Geophys. Res. Lett.*, 13, 1007-1010, 1986.
- Bolt, B. A., and J. A. Canas, Constraints from core reflections on mantle Q and density at the core boundary, *Phys. Earth Planet. Interiors*, 38, 1-8, 1985.
- Bolt, B. A., and L. A. Drake, Love mode dispersion across subduction zones by finite element modelling, *Geophys. J. R. Astron. Soc.*, 84, 515-528, 1986.
- Bolt, B. A., and M. Niaz, S velocities in D* from diffracted SH-waves at the core boundary, *Geophys. J. R. Astron. Soc.*, 79, 825-834, 1984.
- Boss, A. P., and I. S. Sacks, Formation and growth of deep mantle plumes, *Geophys. J. R. Astron. Soc.*, 80, 241-255, 1985.
- Boyd, T. M., and K. C. Creager, Kinematic and travel time constraints on aseismic extensions of the Aleutian slab, *Eos Trans. AGU*, 67, 380, 1986.
- Boyd, T. M., J. A. Snoke, I. S. Sacks, and A. Rodriguez B., High resolution determination of the Benioff zone geometry beneath southern Peru, *Bull. Seismol. Soc. Am.*, 74, 559-568, 1984.
- Budiansky, B., E. E. Sumner, and R. O'Connell, Bulk thermoelastic attenuation of composite materials, *J. Geophys. Res.*, 80, 343-348, 1983.
- Burdick, L. J., Estimation of the frequency dependence of Q from ScP and ScS phases, *Geophys. J. R. Astron. Soc.*, 80, 35-55, 1985.
- Burdick, L. J., and S. P. Grand, Estimation of Q^* for Asian travel paths using SP and S phases, in *The VELA Project*, ed Ann Kerr, 677-692, 1985.
- Burdick, L. J., and C. A. Salavado, Modeling body wave amplitude fluctuations using the three-dimensional slowness method, *J. Geophys. Res.*, 91, 12,482-12,496, 1986.
- Burdick, L. J., T. Wallace, and T. Lay, Modelling the near field and teleseismic observations from the Amchitka test site, *J. Geophys. Res.*, 89, 4373-4388, 1984.
- Burger, R. W., T. Lay, and L. J. Burdick, Average Q and yield estimates from the Pahute Mesa test site, *Bull. Seismol. Soc. Am.*, in press, 1986.
- Busse, F. H., Quadrupole convection in the lower mantle?, *Geophys. Res. Lett.*, 10, 285-288, 1983.
- Butler, R., P wave travel time and amplitude in western North America, *Nature*, 36, 677-678, 1983.
- Butler, R., P wave amplitudes in the United States, east-west variations and variability, *Geophys. J. R. Astron. Soc.*, 77, 767-774, 1984a.
- Butler, R., Azimuth, energy, Q, and temperature variations on P wave amplitudes in the United States, *Rev. Geophys. Space Phys.*, 22, 1-36, 1984b.
- Butler, R., P wave amplitude variation between the eastern and western United States, reply to Z. A. Der, *Geophys. J. R. Astron. Soc.*, 81, 385-388, 1985.
- Butler, R., Amplitudes at the antipode, *Bull. Seism. Soc. Am.*, 76, 1355-1365, 1986.
- Butler, R., T. M. Brocher, and J. A. Rial, Inner core experiments, teleseismic exploration of the antipode, *Eos Trans. AGU*, 67, 89-94, 1986.
- Cahill, T., and B. L. Isacks, An apparent double-planned Benioff zone beneath northern Chile resulting from misidentification of reflected phases, *Geophys. Res. Lett.*, 13, 333-336, 1986.
- Chao, C.-H., T.-L. Teng, and Z. Zhen, The crust and upper mantle of the Pacific Ocean basin, *Eos Trans. AGU*, 64, 759, 1983.
- Chiu, J.-M., B. L. Isacks, and R. K. Cardwell, Propagation of high-frequency seismic waves inside the subducted lithosphere from intermediate-depth earthquakes recorded in the Vanuatu Arc, *J. Geophys. Res.*, 90, 12,741-12,754, 1985.
- Choy, G. L., and V. F. Cormier, The structure of the inner core inferred from short-period and broadband GDSN data, *Geophys. J. R. Astron. Soc.*, 72, 1-21, 1983.
- Choy, G. L., and V. F. Cormier, Direct measurement of the mantle attenuation operator from broadband P and S waveforms, *J. Geophys. Res.*, 91, 7326-7342, 1986.
- Clayton, R. W., and R. P. Comer, A tomographic analysis of mantle heterogeneities from body wave travel time data, *Eos Trans. AGU*, 64, 776, 1983.
- Cockerham, R. S., Evidence for a 180-km-long subducted slab beneath northern California, *Bull. Seismol. Soc. Am.*, 74, 569-576, 1984.
- Comer, R. P., Rapid seismic ray tracing in a spherically symmetric earth via interpolation of rays, *Bull. Seismol. Soc. Am.*, 74, 479-492, 1984.
- Comer, R. P., and R. W. Clayton, Reconstruction of mantle heterogeneity by iterative back-projection of travel times I. theory and reliability, *J. Geophys. Res.*, in press, 1986.
- Cormier, V. F., The effect of attenuation on seismic body waves, *Bull. Seismol. Soc. Am.*, 72, S169-S200, 1982.
- Cormier, V. F., Deep earth structure, *Rev. Geophys. Space Phys.*, 21, 1277-1284, 1983.
- Cormier, V. F., The polarization of S waves in a heterogeneous isotropic earth model, *J. Geophys. Res.*, 86, 20-23, 1984.
- Cormier, V. F., Some problems with S, SKS, and ScS observations and implications for the structure of the base of the mantle and outer core, *J. Geophys. Res.*, 87, 14-22, 1985.
- Cormier, V. F., Synthesis of body waves in transversely isotropic earth models, *Bull. Seismol. Soc. Am.*, 76, 231-240, 1986.
- Cormier, V. F., and G. L. Choy, A search for lateral heterogeneity in the inner core from differential travel times near PKP-D and PKP-C, *Geophys. Res. Lett.*, in press, 1986.
- Creager, K. C., Geometry, velocity structure, and penetration depths of descending slabs in the western Pacific, Ph.D. thesis, University of California, San Diego, 1984.
- Creager, K. C., and T. H. Jordan, Slab penetration into the lower mantle, *J. Geophys. Res.*, 89, 3031-3049, 1984.
- Creager, K. C., and T. H. Jordan, Slab penetration into the lower mantle beneath the Mariana and other island arcs of the northwest Pacific, *J. Geophys. Res.*, 91, 3573-3589, 1986a.
- Creager, K. C., and T. H. Jordan, Aspherical structure of the core-mantle boundary, from PKP travel times, *Geophys. Res. Lett.*, in press, 1986b.
- Crossley, D., Oscillatory flow in the liquid core, *Phys. Earth Planet. Interiors*, 36, 1-16, 1984.
- Cunningham, P. S., S. W. Roecker, and D. Hatzfeld, Three-dimensional P and S wave structures of Southern Peru and their tectonic implications, *J. Geophys. Res.*, 91, 9517-9532, 1986.
- Dahlen, F. A., and I. H. Henson, Asymptotic normal modes of a laterally heterogeneous earth, *J. Geophys. Res.*, 90, 12,653-12,681, 1985.
- Davies, G. F., Geophysical and isotopic constraints on mantle convection an interim synthesis, *J. Geophys. Res.*, 89, 6017-6040, 1984.
- Davis, J. P., Variation in apparent attenuation of the earth's normal modes due to lateral heterogeneity, *Geophys. Res. Lett.*, 12, 141-144, 1985.
- Davis, J. P., Local eigenfrequency and its uncertainty inferred from fundamental spheroidal mode peak shifts, *Eos Trans. AGU*, 67, 306, 1986.
- Davis, J. P., and I. H. Henson, Validity of the great circular average approximation for inversion of normal mode measurements, *Geophys. J. R. Astron. Soc.*, 85, 69-92, 1986.
- Der, Z. A., Comments on 'P wave amplitudes in the United States: east-west variations and variability' by Rhett Butler, *Geophys. J. R. Astron. Soc.*, 81, 381-383, 1985.
- Der, Z. A., and A. C. Lees, Methodologies for estimating $Q^*(f)$ from short-period body waves and regional variations in $Q^*(f)$ in the United States, *Geophys. J. R. Astron. Soc.*, 82, 125-140, 1985.
- Der, Z. A., A. C. Lees, and V. F. Cormier, Frequency dependence of Q in the mantle underlying the shield areas of Eurasia, III, the Q model, *Geophys. J. R. Astron. Soc.*, in press, 1986.
- Der, Z. A., A. C. Lees, L. M. Anderson, J. A. Burnett, M. E. Marshall, T. W. McElfresh, and R. Wagner, Frequency dependence of Q in the mantle under shield areas of Eurasia, in *The VELA Program*, ed Ann Kerr, 693-704, 1985.
- Der, Z. A., T. W. McElfresh, and A. O'Donnell, An investigation of the regional variations and frequency dependence of anelastic attenuation in the mantle under the United States in the 0.5-4.0 Hz band, *Geophys. J. R. Astron. Soc.*, 69, 67-99, 1982a.
- Der, Z. A., W. D. Rivers, T. W. McElfresh, A. O'Donnell, P. J. Klouda, and M. E. Marshall, Worldwide variations in the attenuative properties of the upper mantle as determined by spectral studies of short period body waves, *Phys. Earth Planet. Interiors*, 30, 12-25, 1982b.
- Dziewonski, A. M., Mapping the lower mantle: determination of lateral heterogeneity in P velocity up to degree and order 6, *J. Geophys. Res.*, 89, 5929-5952, 1984.
- Dziewonski, A. M., and D. L. Anderson, Travel times and station corrections for P waves at teleseismic distances, *J. Geophys. Res.*, 88, 3295-3314, 1983.
- Dziewonski, A. M., and D. L. Anderson, Seismic tomography of the earth's interior, *Am. Sci.*, 721, 483-494, 1984.
- Dziewonski, A. M., and J. M. Steim, Dispersion and attenuation of mantle waves through waveform inversion, *Geophys. J. R. Astron. Soc.*, 70, 503-527, 1982.
- Estey, L. H., and B. J. Douglas, Upper mantle anisotropy: A preliminary model, *J. Geophys. Res.*, 91, 11,393-11,406, 1986.
- Feng, C.-C., and T.-L. Teng, Three-dimensional crust and upper mantle structure of the Eurasian continent, *J. Geophys. Res.*, 88, 2261-2272, 1983.
- Fischer, K. M., K. C. Creager, and T. H. Jordan, Mapping the Tonga slab, *Eos Trans. AGU*, 67, 381, 1986.
- Frankel, A., and R. W. Clayton, A finite difference simulation of wave propagation in two-dimensional random media, *Bull. Seismol. Soc. Am.*, 74, 2167-2186, 1984.
- Friedlander, S., Internal oscillations in the earth's fluid core, *Geophys. J. R. Astron. Soc.*, 80, 345-361, 1985.
- Gee, L. S., and T. H. Jordan, Earth structure from hybrid representations of seismograms, *Eos Trans. AGU*, 67, 302, 1986.
- Giardini, D., and J. H. Woodhouse, Deep seismicity and modes of deformation in Tonga subduction zone, *Nature*, 307, 505-509, 1984.
- Giardini, D., X.-D. Li, and J. H. Woodhouse, Three dimensional structure of the earth from splitting in free oscillation spectra, *Nature*, in press, 1986.
- Given, J. W., Inversion of body-wave seismograms for upper mantle structure, Ph.D. thesis, 154 pp., Calif. Inst. of Technol., Pasadena, 1983.
- Grand, S. P., Shear velocity structure of the mantle beneath the North American plate, Ph.D. thesis, 228 pp., Calif. Inst. of Technol., Pasadena, 1986a.
- Grand, S. P., A tomographic inversion for shear velocity beneath the North American plate, *Eos Trans. AGU*, 67, 302, 1986b.
- Grand, S. P., and D. V. Helmerger, Upper mantle shear structure of North America, *Geophys. J. R. Astron. Soc.*, 76, 399-438, 1984a.
- Grand, S. P., and D. V. Helmerger, Upper mantle shear structure beneath the northwest Atlantic ocean, *J. Geophys. Res.*, 89, 11,465-11,475, 1984b.
- Grand, S. P., and D. V. Helmerger, Upper mantle shear structure beneath Asia from multi-bounce S waves, *Phys. Earth Planet. Interiors*, 41, 154-169, 1985.
- Grange, F., P. Cunningham, J. Gagnepain, D. Hatzfeld, P. Molnar, L. Ocola, A. Rodriguez, S. W. Roecker, J. M. Stock, and G. Suarez, The configuration of the seismic zone and the downgoing slab in southern Peru, *Geophys. Res. Lett.*, 11, 38-41, 1984.
- Graves, R. S., Grand, and D. V. Helmerger, Upper mantle cross-section from Tonga to California, *Eos Trans. AGU*, 66, 975, 1985.
- Hager, B. H., Subducted slabs and the geoid: constraints on mantle rheology and flow, *J. Geophys. Res.*, 89, 6003-6015, 1984.
- Hager, B. H., R. W. Clayton, M. A. Richards, R. P. Comer, and A. M. Dziewonski, Lower mantle heterogeneity, dynamic topography and the geoid, *Nature*, 313, 541-545, 1985.
- Hansen, R. A., Simultaneous estimation of terrestrial eigenfunctions, *Geophys. J. R. Astron. Soc.*, 70, 155-172, 1982.
- Hauksson, E., Structure of the Benioff zone beneath the Shumagin Islands, Alaska: relocation of local earthquakes using three-dimensional ray tracing, *J. Geophys. Res.*, 90, 635-649, 1985.
- Helmerger, D. V., Theory and application of synthetic seismograms, *Proceedings of the Enrico Fermi International School of Physics, LXXXV*, eds H. Kanamori and E. Bosch, 174-222, 1983.
- Helmerger, D. V., G. Engen, and S. Grand, Notes on wave propagation in laterally varying structure, *J. Geophys. Res.*, 88, 82-91, 1985a.
- Helmerger, D. V., G. Engen, and S. Grand, Upper-mantle cross-section from California to Greenland, *J. Geophys. Res.*, 88, 92-100, 1985b.
- Henson, I. H., and F. A. Dahlen, Asymptotic normal modes of a laterally heterogeneous earth 2. Further results, *J. Geophys. Res.*, 91, 12,467-12,481, 1986.
- Humphreys, E. R., W. Clayton, and B. H. Hager, A tomographic image of mantle structure beneath southern California, *Geophys. Res. Lett.*, 11, 625-627, 1984.
- Incorporated Research Institutions for Seismology, Science Plan for a New Global Seismographic Network, Washington, D. C., 1984.
- Ingate, S. F., G. L. Duckworth, K. J. Muirhead, and V. Cormier, Inversion of high-resolution slant-stack data for upper mantle structure, *Eos Trans. AGU*, 66, 974, 1985.
- Iyer, H. M., A review of crust and upper mantle structure studies of the Snake River plain-Yellowstone volcanic system: a major lithospheric anomaly in the western U.S.A., *Tectonophysics*, 105, 291-308, 1984.
- Jeanloz, R., and A. B. Thompson, Phase transitions and mantle discontinuities, *Rev. Geophys. Space Phys.*, 21, 51-74, 1983.
- Johnson, L. R., and R. C. Lee, Extremal bounds on the P-velocity in the earth's core, *Bull. Seismol. Soc. Am.*, 75, 115-130, 1985.
- Jordan, T. H., and K. C. Creager, Chemical boundary layers and large-scale flow in the mantle and core, *Eos Trans. AGU*, 67, 311, 1986.
- Jordan, T. H., A. L. Lerner-Lam, and K. C. Creager, Seismic imaging of mantle convection: the evidence for deep circulation, in *Mantle Convection*, W. R. Pelletier (ed.), in press, 1986.
- Kawakatsu, H., Can "pure path" models explain free oscillation data?, *Geophys. Res. Lett.*, 10, 186-189, 1983.
- Kawakatsu, H., Double seismic zone in Tonga, *Nature*, 316, 53-55, 1985.
- Kawakatsu, H., Double seismic zones kinematics, *J. Geophys. Res.*, 91, 4811-4825, 1986a.
- Kawakatsu, H., Downward-tensional earthquakes beneath the Tonga arc: A double seismic zone?, *J. Geophys. Res.*, 91, 6432-6440, 1986b.
- Kawakatsu, H., and T. Seno, Triple seismic zone and the regional variation of seismicity along the northern Honshu arc, *J. Geophys. Res.*, 88, 4215-4230, 1983.
- Kawasaki, I., A method for the near-source anisotropy by the pair-event inversion of Rayleigh-wave radiation patterns, *Geophys. J. R. Astron. Soc.*, 71, 395-424, 1982.
- Knopoff, L., The thickness of the lithosphere from the dispersion of surface waves, *Geophys. J. R. Astron. Soc.*, 74, 55-81, 1983.
- Kuo, B.-Y., D. W. Forsyth, and M. Wyssession, Lateral heterogeneity and azimuthal anisotropy in the north Atlantic determined from SS-S differential travel times, *J. Geophys. Res.*, in press, 1986.
- Langston, C. A., and W. A. Arnold, Moment tensor inverses from dipping slabs, *Geophys. Res. Lett.*, 9, 1290-1293, 1982.
- Langston, C. A., and C.-E. Baag, The validity of ray theory approximations for the computation of teleseismic SV waves, *Bull. Seismol. Soc. Am.*, 75, 1719-1727, 1985.
- Lavelle, E. M., D. W. Forsyth, and P. Friedemann, Scales of heterogeneity near the core-mantle boundary, *Geophys. Res. Lett.*, in press, 1986.
- Lay, T., Localized velocity anomalies in the lower mantle, *Geophys. J. R. Astron. Soc.*, 72, 483-516, 1983.
- Lay, T., Analysis of diffracted S waves traversing a layer with a lower mantle shear velocity discontinuity, *Eos Trans. AGU*, 66, 310, 1985.
- Lay, T., Investigations of D* shear velocity structure using the Graefenberg array, *Eos Trans. AGU*, 67, 312, 1986a.
- Lay, T., Evidence for a lower mantle shear velocity discontinuity in S and sS phases, *Geophys. Res. Lett.*, in press, 1986b.
- Lay, T., and D. V. Helmerger, Body-wave amplitude and travel time correlations across North America, *Bull. Seismol. Soc. Am.*, 73, 1063-1076, 1983a.
- Lay, T., and D. V. Helmerger, A lower mantle S wave triplication and the shear velocity structure of D*, *Geophys. J. R. Astron. Soc.*, 75, 799-837, 1983b.
- Lay, T., and D. V. Helmerger, A shear velocity discontinuity in the lower mantle, *Geophys. Res. Lett.*, 10, 63-66, 1983c.
- Lay, T., and D. V. Helmerger, The shear-wave velocity gradient at the base of the mantle, *J. Geophys. Res.*, 88, 8160-8170, 1983d.
- Lay, T., and H. Kanamori, Geometric effects of global lateral heterogeneity on long-period surface wave propagation, *J. Geophys. Res.*, 90, 605-621, 1985.
- Lay, T., and T. C. Wallace, Multiple ScS travel times and attenuation beneath Mexico and Central America, *Geophys. Res. Lett.*, 10, 301-304, 1983.
- Lay, T., and C. J. Young, The effect of SKS scattering on models of the shear-velocity structure of the D* region, *J. Geophys. Res.*, 91, 11-15, 1986.
- Lee, R. C., and L. R. Johnson, Tau estimates for mantle P and S waves from global travel time observations, *Geophys. J. R. Astron. Soc.*, 77, 655-666, 1984a.
- Lee, R. C., and L. R. Johnson, Extremal bounds on the seismic velocities in the earth's mantle, *Geophys. J. R. Astron. Soc.*, 77, 667-681, 1984b.
- Lees, A. C., M. S. T. Bukowski, and R. Jeanloz, Reflection properties of phase transition and compositional change models of the 670-km discontinuity, *J. Geophys. Res.*, 88, 8145-8159, 1983.
- Lefevre, L. V., and D. V. Helmerger, Variations in upper mantle P wave velocity beneath North America, *Eos Trans. AGU*, 65, 234, 1984.
- Lerner-Lam, A. L., The structure of the oceanic lithosphere from anisotropic inversion of surface and body waveforms, *Eos Trans. AGU*, 67, 302, 1986.
- Lerner-Lam, A. L., and T. H. Jordan, Earth structure from fundamental and higher-mode waveform analysis, *Geophys. J. R. Astron. Soc.*, 75, 759-797, 1983.
- Lerner-Lam, A. L., and T. H. Jordan, How thick are the continents?, *J. Geophys. Res.*, in press, 1986.
- Loper, D. E., The dynamical structures of D* and deep plumes in a non-

- Newtonian mantle. *Phys Earth Planet Interiors*, 34, 57-67, 1984.
- Loper, D. E., and D. R. Fearn, A seismic model of a partially molten inner core. *J. Geophys. Res.*, 88, 1235-1242, 1983.
- Loper, D. E., and F. D. Stacey, The dynamical and thermal structure of deep mantle plumes. *Phys Earth Planet Interiors*, 33, 304-317, 1983.
- Lynnes, C., and T. Lay, Decoupling of short period P waves by a high velocity anomaly beneath Pahute Mesa. *Eos Trans. AGU*, 65, 994, 1984.
- Masters, G., and F. Gilbert, Attenuation in the earth at low frequencies. *Philos. Trans. R. Soc. London Ser. A*, 308, 479-522, 1983.
- Masters, G., and F. Gilbert, Spheroidally averaged structure from free oscillation data. *Eos Trans. AGU*, 66, 966, 1985.
- Masters, G., T. H. Jordan, P. G. Silver, and F. Gilbert, Aspherical earth structure from fundamental spheroidal-mode data. *Nature*, 298, 609-613, 1982.
- Masters, G., J. Park, and F. Gilbert, Observations of coupled spheroidal and toroidal modes. *J. Geophys. Res.*, 88, 10,285-10,298, 1983.
- Mavko, B. B., and G. A. Thompson, Crustal and upper mantle structure of the northern and central Sierra Nevada. *J. Geophys. Res.*, 88, 5874-5892, 1983.
- Menke, W., A formula for the apparent attenuation of acoustic waves in randomly layered media. *Geophys. J. R. Astron. Soc.*, 75, 541-544, 1983.
- Menke, W., Effect of heterogeneities in D'' on the decay rate of Pd_{iff} . *J. Geophys. Res.*, 91, 1927-1933, 1986a.
- Menke, W., Few 2-50 km corrugations on the core-mantle boundary. *Geophys. Res. Lett.*, in press, 1986b.
- Michaelson, C. A., and C. S. Weaver, Upper mantle structure from teleseismic P wave arrivals in Washington and northern Oregon. *J. Geophys. Res.*, 91, 2077-2094, 1986.
- Mitchell, B. J., On the inversion of Love and Rayleigh wave dispersion and implications for earth structure and anisotropy. *Geophys. J. R. Astron. Soc.*, 76, 233-241, 1984.
- Molnar, P., and W.-P. Chen, S-P wave travel time residuals and lateral inhomogeneity in the mantle beneath Tibet and the Himalaya. *J. Geophys. Res.*, 89, 6911-6917, 1984.
- Morelli, A., A. M. Dziewonski, and J. H. Woodhouse, Anisotropy of the inner core inferred from PKIKP travel times. *Geophys. Res. Lett.*, in press, 1986.
- Morris, S. P., and R. J. Geller, Toroidal modes of a simple laterally heterogeneous sphere. *Bull. Seismol. Soc. Am.*, 72, 1155-1166, 1982.
- Murtha, P. E., Seismic velocities in the upper part of the earth's core. Ph.D. thesis, 115 pp., Univ. Calif. Berkeley, 1984.
- Murtha, P. E., and T. Tanimoto, An investigation of the 650 km discontinuity using Pd_{iff} reflection/transmission coefficients. *Eos Trans. AGU*, 63, 1035, 1982.
- Nakanishi, I., and D. L. Anderson, World-wide distribution of group velocity of mantle Rayleigh waves as determined by spherical harmonic inversion. *Bull. Seismol. Soc. Am.*, 72, 1185-1194, 1982.
- Nakanishi, I., and D. L. Anderson, Measurement of mantle wave velocities and inversion for lateral heterogeneity and anisotropy I. Analysis of great circle phase velocities. *J. Geophys. Res.*, 88, 10,267-10,283, 1983.
- Nakanishi, I., and D. L. Anderson, Aspherical heterogeneity of the mantle from phase velocities of mantle waves. *Nature*, 307, 117-121, 1984a.
- Nakanishi, I., and D. L. Anderson, Measurements of mantle wave velocities and inversion for lateral heterogeneity and anisotropy. 2. Analysis by the single-station method. *Geophys. J. R. Astron. Soc.*, 78, 573-617, 1984b.
- Nakanishi, I., and H. Kanamori, Effects of lateral heterogeneity and source process time on the linear moment tensor inversion of long-period Rayleigh waves. *Bull. Seismol. Soc. Am.*, 72, 2063-2080, 1982.
- Nataf, H.-C., I. Nakanishi, and D. L. Anderson, Anisotropy and shear-velocity heterogeneities in the upper mantle. *Geophys. Res. Lett.*, 11, 109-112, 1984.
- Nataf, H.-C., I. Nakanishi, and D. L. Anderson, Measurements of mantle wave velocities and inversion for lateral heterogeneities and anisotropy 3. Inversion. *J. Geophys. Res.*, 91, 7261-7307, 1986.
- Nishimura, C. E., and D. W. Forsyth, Anomalous Love-wave phase velocities in the Pacific: sequential pure-path and spherical harmonic inversion. *Geophys. J. R. Astron. Soc.*, 81, 389-407, 1985.
- Nishimura, C., and D. W. Forsyth, Rayleigh wave phase velocities in the Pacific with implications for azimuthal anisotropy and lateral heterogeneities. *Geophys. J. R. Astron. Soc.*, in press, 1986.
- Okal, E. A., Attenuation measurements for mantle Rayleigh spheroidal overtones (3R, 4R, 5R). *Eos Trans. AGU*, 67, 307, 1986.
- Okal, E. A., and B.-G. Jo, Regional dispersion of first-order overtone Rayleigh waves. *Geophys. J. R. Astron. Soc.*, 72, 461-481, 1983.
- Okal, E. A., and B.-G. Jo, Stacking investigations of higher-order mantle Rayleigh waves. *Geophys. Res. Lett.*, 12, 421-424, 1985.
- Park, J., Synthetic seismograms from coupled free oscillations: effects of lateral structure and rotation. *J. Geophys. Res.*, 91, 6441-6464, 1986.
- Park, J., and F. Gilbert, Coupled free oscillations of an aspherical, dissipative, rotating earth. Galerkin theory. *J. Geophys. Res.*, 91, 7241-7260, 1986.
- Patton, H. J., and S. R. Taylor, Q structure of the Basin and Range from surface waves. *J. Geophys. Res.*, 89, 6929-6940, 1984.
- Place, D. J., and J. E. Ebel, Array measurements for the lower mantle from the New England seismic network. *Eos Trans. AGU*, 63, 1035, 1982.
- Powell, C. A., and A. S. Meltzer, Scattering of P waves beneath SCARLET in southern California. *Geophys. Res. Lett.*, 11, 481-484, 1984.
- Priestley, K., and D. E. Chavez, Magnitude bias in the Great Basin and its implications for explosion magnitude versus yield estimates. *Geophys. Res. Lett.*, 12, 573-576, 1985.
- Regan, J., and D. L. Anderson, Anisotropic models of the upper mantle. *Phys Earth Planet Interiors*, 35, 227-263, 1984.
- Revenaugh, J. S., and T. H. Jordan, Observation of first-order mantle reverberations. *Eos Trans. AGU*, 66, 967, 1985.
- Rial, J. A., S. Grand, and D. V. Helmberger, A note on lateral variation in upper mantle shear-wave velocity across the Alpine front. *Geophys. J. R. Astron. Soc.*, 77, 639-654, 1984.
- Richards, M. A., and B. H. Hager, Geoid anomalies in a dynamic earth. *J. Geophys. Res.*, 89, 5987-6002, 1984.
- Richards, P. G., Seismic wave propagation effects - development of theory and numerical modeling, in *The VELA Program*, ed. Ann Kerr, 183-226, 1985.
- Richards, P. G., and W. Menke, The apparent attenuation of a scattering medium. *Bull. Seismol. Soc. Am.*, 73, 1005-1021, 1983.
- Ritzwiler, M. G., Masters, G., and F. Gilbert, Observations of anomalous splitting and their interpretation in terms of aspherical structure. *J. Geophys. Res.*, 91, 10,203-10,228, 1986.
- Roecker, S. W., Velocity structure in the Izu-Bonin seismic zone and the depth of the olivine-spinel phase transition in the slab. *J. Geophys. Res.*, 90, 7771-7794, 1985.
- Rohay, A. C., Control of subducted slab geometry beneath the Washington Cascades by continental deformation. *Eos Trans. AGU*, 63, 1023, 1982.
- Rosa, J. W. C., and K. Aki, Regionalized Rayleigh wave phase and group velocity. *Eos Trans. AGU*, 67, 306-307, 1986.
- Ruff, L., and E. Lettvin, Short period P wave amplitudes and the variability of the core shadow zone. *Eos Trans. AGU*, 65, 999, 1984.
- Rytov, P. A., and L. Knopoff, Spectral analysis of gapped data search for ISI of the south pole. *J. Geophys. Res.*, 89, 1899-1902, 1984.
- Schwartz, S., and T. Lay, Comparison of long-period surface wave amplitude and phase anomalies for two models of global lateral heterogeneity. *Geophys. Res. Lett.*, 12, 231-234, 1985.
- Shore, M. J., A seismic Q-profile for the lower mantle from short period P waves. *Nature*, 310, 399-401, 1984.
- Silver, P. G., and W. W. Chan, Observations of body wave multipathing from broadband seismograms: evidence for lower-mantle slab penetration beneath the Sea of Okhotsk. *J. Geophys. Res.*, in press, 1986.
- Snoke, J. A., and I. S. Sacks, Seismic modeling of lateral heterogeneity at the base of the mantle. *Geophys. J. R. Astron. Soc.*, 86, 801-814, 1986.
- Souriau, A., On the retrieval of pure path velocities from great circle data. *Geophys. J. R. Astron. Soc.*, 80, 783-790, 1985.
- Souriau, A., and J. H. Woodhouse, A worldwide comparison of predicted S-wave delays from a three dimensional upper mantle model with P-wave station corrections. *Phys. Earth Planet. Interiors*, 39, 75-88, 1985.
- Spencer, C. P., and E. R. Engdahl, A joint hypocentre location and velocity inversion technique applied to the central Aleutians. *Geophys. J. R. Astron. Soc.*, 72, 399-415, 1983.
- Stacey, F. D., and D. E. Loper, The thermal boundary layer interpretation of D'' and its role as a plume source. *Phys. Earth Planet. Interiors*, 33, 45-55, 1983.
- Stark, M., and D. W. Forsyth, The geoid, small-scale convection, and differential travel time anomalies of shear waves in the central Indian ocean. *J. Geophys. Res.*, 88, 2273-2288, 1983.
- Stark, P. B., and C. Frohlich, The depths of the deepest deep earthquakes. *J. Geophys. Res.*, 90, 1859-1869, 1985.
- Stark, P. B., I. Parker, G. Masters, and J. A. Orcutt, Travel time constraints on core structure. *Eos Trans. AGU*, 67, 311, 1986.
- Stork, C., and D. V. Helmberger, A new velocity profile for the Eurasian shield. *Eos Trans. AGU*, 66, 310, 1985.
- Stefani, J. P., R. J. Geller, and G. C. Kroeger, A direct measurement of the distance between a hypocenter in a Benioff-Wadati zone and the slab-aesthenosphere contact. *J. Geophys. Res.*, 87, 323-328, 1982.
- Suyehiro, K., and I. S. Sacks, An anomalous low velocity region above the deep earthquakes in the Japan subduction zone. *J. Geophys. Res.*, 88, 10,429-10,438, 1985.
- Taber, J. J., and K. W. Hudnut, A transition from a single to a double Benioff zone near the Shumagin Islands, Alaska. *Eos Trans. AGU*, 66, 958, 1985.
- Tajima, F., and J. D. Garmany, Surface wave amplitude asymmetries: is reciprocity valid? *Phys. Earth Planet. Interiors*, in press, 1986.
- Tanimoto, T., A simple derivation of the formula to calculate synthetic long-period seismograms in a heterogeneous earth by normal mode summation. *Geophys. J. R. Astron. Soc.*, 77, 275-278, 1984a.
- Tanimoto, T., Waveform inversion of mantle Love waves: the Born seismogram approach. *Geophys. J. R. Astron. Soc.*, 78, 641-660, 1984b.
- Tanimoto, T., The Backus-Gilbert approach to the three-dimensional structure in the upper mantle. I. Lateral variation of surface wave phase velocity with its error and resolution. *Geophys. J. R. Astron. Soc.*, 82, 105-124, 1985.
- Tanimoto, T., Free oscillations in a slightly anisotropic earth. *Geophys. J. R. Astron. Soc.*, 87, 493-517, 1986a.
- Tanimoto, T., The Backus-Gilbert approach to the three-dimensional structure in the upper mantle. II, SH and SV velocity. *Geophys. J. R. Astron. Soc.*, 84, 49-70, 1986b.
- Tanimoto, T., and D. L. Anderson, Mapping convection in the mantle. *Geophys. Res. Lett.*, 11, 287-290, 1984.
- Tanimoto, T., and D. L. Anderson, Lateral heterogeneity and azimuthal anisotropy of the upper mantle: Love and Rayleigh waves 100-250 s. *J. Geophys. Res.*, 90, 1842-1858, 1985.
- Tanimoto, T., and B. A. Bolt, Coupling of torsional modes in the earth. *Geophys. J. R. Astron. Soc.*, 74, 83-96, 1983.
- Tanimoto, T., and H. Kanamori, Linear programming approach to moment tensor inversion of earthquake sources and some tests on the three dimensional structure of the upper mantle. *Geophys. J. R. Astron. Soc.*, 84, 413-430, 1986.
- Taylor, S. R., Three dimensional crust and upper mantle structure at the Nevada test site. *J. Geophys. Res.*, 88, 2220-2232, 1983.
- Taylor, S. R., B. P. Bonner, and G. Zandt, Attenuation and scattering of broadband P and S waves across North America. *J. Geophys. Res.*, 91, 7309-7325, 1986.
- Tralli, D. M., and L. R. Johnson, Estimation of slowness-dependent source and receiver corrections for P wave travel times. *Bull. Seism. Soc. Am.*, 76, 1718-1738, 1986a.
- Tralli, D. M., and L. R. Johnson, Lateral variations in mantle P velocity from tectonically regionalized tau estimates. *Geophys. J. R. Astron. Soc.*, 86, 475-489, 1986.
- Tsuboi, S., R. J. Geller, and S. P. Morris, Partial derivatives of the eigenfrequencies of a laterally heterogeneous earth model. *Geophys. Res. Lett.*, 12, 817-820, 1985.
- Vassiliou, M. S., The state of stress in subducting slabs as revealed by earthquakes analyzed by moment tensor inversion. *Earth Planet. Sci. Lett.*, 69, 195-202, 1984.
- Vassiliou, M. S., B. H. Hager, and A. Raefsky, The distribution of earthquakes with depth and stress in subducting slabs. *J. Geodyn.*, 1, 11-28, 1984.
- Walck, M. C., The P wave upper mantle structure beneath an active spreading centre: the Gulf of California. *Geophys. J. R. Astron. Soc.*, 76, 697-723, 1984.
- Walck, M. C., The upper mantle beneath the northeast Pacific rim: a comparison with the Gulf of California. *Geophys. J. R. Astron. Soc.*, 81, 243-276, 1985.
- Walck, M. C., and R. W. Clayton, Analysis of upper mantle structure using wavefield continuation of P waves. *Bull. Seismol. Soc. Am.*, 74, 1703-1719, 1984.
- Wielandt, E., and L. Knopoff, Dispersion of very long-period Rayleigh waves along the East Pacific Rise: evidence for S wave velocity anomalies to 450 km depth. *J. Geophys. Res.*, 87, 8631-8641, 1982.
- Wong, Y. K., and J. H. Woodhouse, Ray theory for surface waves on a sphere. *Eos Trans. AGU*, 64, 260, 1983.
- Wong, Y. K., and J. H. Woodhouse, The mapping of mantle wave phase velocity in the presence of large deviations in ray-path. *Eos Trans. AGU*, 67, 307, 1986.
- Woodhouse, J. H., The joint inversion of seismic waveforms for lateral variations in earth structure and earthquake source parameters. *Proceedings of the Enrico Fermi International School of Physics, LXXXV*, eds H. Kanamori and E. Boschi, 366-397, 1984.
- Woodhouse, J. H., and A. M. Dziewonski, Mapping the upper mantle three-dimensional modeling of earth structure by inversion of seismic waveforms. *J. Geophys. Res.*, 89, 5953-5986, 1984.
- Woodhouse, J. H., and A. M. Dziewonski, Three-dimensional images of the earth's interior. in *The VELA Program*, ed. Ann Kerr, 448-464, 1985.
- Woodhouse, J. H., and A. M. Dziewonski, Three dimensional mantle models based on mantle wave and long period body wave data. *Eos Trans. AGU*, 67, 307, 1986.
- Woodhouse, J. H., and T. P. Gurnis, Surface waves and free oscillations in a regionalized earth model. *Geophys. J. R. Astron. Soc.*, 88, 653-673, 1982.
- Woodhouse, J. H., D. Giardini, and X.-D. Li, Evidence for inner core anisotropy from splitting in free oscillation spectra. *Geophys. Res. Lett.*, in press, 1986.
- Yomogida, K., Gaussian beams for surface waves in laterally slowly varying media. *Geophys. J. R. Astron. Soc.*, 82, 511-533, 1985.
- Yomogida, K., and K. Aki, Waveform synthesis of surface waves in a laterally heterogeneous earth by the Gaussian beam method. *J. Geophys. Res.*, 90, 7665-7688, 1985.
- Yomogida, K., and K. Aki, Amplitude and phase data inversions for phase velocity anomalies in the Pacific ocean basin. *Geophys. J. R. Astron. Soc.*, in press, 1986.
- Young, C. J., and T. Lay, Evidence for a shear velocity discontinuity in the lowermost mantle beneath India and the Indian ocean. *Eos Trans. AGU*, 67, 311-312, 1986a.
- Young, C. J., and T. Lay, The core-mantle boundary. *Annual Rev. Earth Planet. Sci.*, in press, 1986b.
- Zandt, G., and G. E. Randall, Observations of shear-coupled P waves. *Geophys. Res. Lett.*, 12, 565-568, 1985.
- Zhang, J., and T. Lay, Investigation of a lower mantle shear wave triplication using a broadband array. *Geophys. Res. Lett.*, 11, 620-623, 1984.
- Zhou, H.-W., and R. W. Clayton, Mantle velocities from PP waves. *Eos Trans. AGU*, 66, 975, 1985.

(Received September 22, 1986;
accepted December 17, 1986.)